

УДК 530.145

© Кауффман Л. Х., 2022

**ПРИМЕЧАНИЯ К УРАВНЕНИЮ ДИРАКА**Кауффман Л. Х.<sup>a,1</sup>

<sup>a</sup> Иллинойский университет в Чикаго, 851 Саут Морган Стрит, Чикаго, Иллинойс, 60607-7045, США.

В данной работе рассматривается структура уравнения Дирака и дается новая трактовка уравнения Дирака в 1+1 пространстве - времени.

*Ключевые слова:* Алгебра Клиффорда, уравнение Дирака, шахматная доска Фейнмана.

**REMARKS ON THE DIRAC EQUATION**Kauffman L. H.<sup>a,1</sup>

<sup>a</sup> University of Illinois at Chicago, 851 South Morgan Street, Chicago, Illinois, 60607-7045, USA.

This paper examines the structure of the Dirac equation and gives a new treatment of the Dirac equation in 1+1 spacetime.

*Keywords:* Clifford algebra, Dirac Equation, Feynman checkerboard.

PACS: 03.65.Pm

DOI: 10.17238/issn2226-8812.2022.3.64-73

**Introduction**

This paper is a discussion of the structure of the Dirac equation. Section 2 discusses how the nilpotent and Majorana operators arise naturally. This section provides a link between our work and the work of Peter Rowlands [10]. We end this section with an expression in split quaternions for the Majorana Dirac equation in one dimension of time and three dimensions of space. The Majorana Dirac equation can be written as follows:

$$(\partial/\partial t + \hat{\eta}\eta\partial/\partial x + \epsilon\partial/\partial y + \hat{\epsilon}\eta\partial/\partial z - \hat{\epsilon}\hat{\eta}\eta m)\psi = 0$$

where  $\eta$  and  $\epsilon$  are the simplest generators of iterant algebra with  $\eta^2 = \epsilon^2 = 1$  and  $\eta\epsilon + \epsilon\eta = 0$ , and  $\hat{\epsilon}, \hat{\eta}$  form a copy of this algebra that commutes with it. This combination of the simplest Clifford algebra with itself is the underlying structure of Majorana Fermions, forming indeed the underlying structure of all Fermions. In Section 3 we apply our methods to the Majorana Dirac Equation and give actual real solutions to the equation. These solutions inevitably make direct use of the Majorana Fermion Clifford algebra. This shows explicitly that Fermions and Majorana Fermions are related by the algebraic transformation between Fermion and Clifford algebra. In Section 4, we end the paper in one dimension of space and one dimension of time. We reformulate the Dirac equation for 1+1 spacetime in general, and show that it is related to a Fermionic Clifford algebra that is described in this section. Specializing to a representation of this algebra we give explicit solutions to the Dirac equation and discuss their relationship with the Feynman checkerboard [1, 2].

---

<sup>1</sup>E-mail: kauffman@uic.edu

## 1. The Dirac Equation and Majorana Fermions

We construct the Dirac equation. The algebra underlying this equation has the same properties as the creation and annihilation algebra for fermions. It is by way of this algebra that we will come to the Dirac equation. If the speed of light is equal to 1 (by convention), then energy  $E$ , momentum  $p$  and mass  $m$  are related by the (Einstein) equation

$$E^2 = p^2 + m^2.$$

Dirac constructed his equation by looking for an algebraic square root of  $p^2 + m^2$  so that he could have a linear operator for  $E$  that would take the same role as the Hamiltonian in the Schrödinger equation. We will get to this operator by first taking the case where  $p$  is a scalar (we use one dimension of space and one dimension of time.). Let  $E = \alpha p + \beta m$  where  $\alpha$  and  $\beta$  are elements of a possibly non-commutative, associative algebra. Then

$$E^2 = \alpha^2 p^2 + \beta^2 m^2 + pm(\alpha\beta + \beta\alpha).$$

Hence we will satisfy  $E^2 = p^2 + m^2$  if  $\alpha^2 = \beta^2 = 1$  and  $\alpha\beta + \beta\alpha = 0$ . This is a Clifford algebra pattern.

We have the Dirac equation  $\hat{E} = \alpha\hat{p} + \beta\hat{m}$ . Because the quantum operator for momentum is  $\hat{p} = -i\partial/\partial x$ , the operator for energy is  $\hat{E} = i\partial/\partial t$ , and the operator for mass is  $\hat{m} = m$ , the Dirac equation becomes the differential equation below.

$$i\partial\psi/\partial t = -i\alpha\partial\psi/\partial x + \beta m\psi.$$

Let  $\mathcal{O} = i\partial/\partial t + i\alpha\partial/\partial x - \beta m$  so that the Dirac equation takes the form  $\mathcal{O}\psi(x, t) = 0$ .

Note that  $\mathcal{O}e^{i(px-Et)} = (E - \alpha p - \beta m)e^{i(px-Et)}$ .

We let  $\Delta = (E - \alpha p - \beta m)$  and let  $U = \Delta\beta\alpha = (E - \alpha p - \beta m)\beta\alpha = \beta\alpha E + \beta p - \alpha m$ , so that  $U^2 = -E^2 + p^2 + m^2 = 0$ .

This nilpotent element leads to a (plane wave) solution to the Dirac equation as follows: We have shown that  $\mathcal{O}\psi = \Delta\psi$  for  $\psi = e^{i(px-Et)}$ . It then follows that  $\mathcal{O}(\beta\alpha\Delta\beta\alpha\psi) = \Delta\beta\alpha\Delta\beta\alpha\psi = U^2\psi = 0$ , from which it follows that  $\psi = Ue^{i(px-Et)}$  is a (plane wave) solution to the Dirac equation.

This calculation suggests that we should multiply the operator  $\mathcal{O}$  by  $\beta\alpha$  on the right, obtaining the operator

$$\mathcal{D} = \mathcal{O}\beta\alpha = i\beta\alpha\partial/\partial t + i\beta\partial/\partial x - \alpha m,$$

and the equivalent Dirac equation  $\mathcal{D}\psi = 0$ . For the specific  $\psi$  above we will now have  $\mathcal{D}(Ue^{i(px-Et)}) = U^2e^{i(px-Et)} = 0$ . This idea for reconfiguring the Dirac equation in relation to nilpotent algebra elements  $U$  is due to Peter Rowlands [10]. Rowlands does this in the context of quaternion algebra. Note that the solution to the Dirac equation that we have found is expressed in Clifford algebra. It can be articulated into specific vector solutions by using a matrix representation of the algebra.

We see that  $U = \beta\alpha E + \beta p - \alpha m$  with  $U^2 = 0$  is really the essence of this plane wave solution to the Dirac equation. This means that a natural non-commutative algebra arises directly and can be regarded as the essential information in a Fermion. It is natural to compare this algebra structure with algebra of creation and annihilation operators that occur in quantum field theory.

If we let  $\tilde{\psi} = e^{i(px+Et)}$  (reversing time), then we have  $\mathcal{D}\tilde{\psi} = (-\beta\alpha E + \beta p - \alpha m)\tilde{\psi} = U^\dagger\tilde{\psi}$ , giving a definition of  $U^\dagger$  corresponding to the anti-particle for  $U\psi$ .

We have  $U = \beta\alpha E + \beta p - \alpha m$  and  $U^\dagger = -\beta\alpha E + \beta p - \alpha m$ .

Note that here we have  $(U + U^\dagger)^2 = (2\beta p + \alpha m)^2 = 4(p^2 + m^2) = 4E^2$ , and  $(U - U^\dagger)^2 = -(2\beta\alpha E)^2 = -4E^2$ .

We have that  $U^2 = (U^\dagger)^2 = 0$  and  $UU^\dagger + U^\dagger U = 4E^2$ . Thus we have a direct appearance of the Fermion algebra corresponding to the Fermion plane wave solutions to the Dirac equation. Furthermore, the decomposition of  $U$  and  $U^\dagger$  into the corresponding Majorana Fermion operators corresponds to  $E^2 = p^2 + m^2$ .

Normalizing by dividing by  $2E$  we have  $A = (\beta p + \alpha m)/E$  and  $B = i\beta\alpha$ . so that  $A^2 = B^2 = 1$  and  $AB + BA = 0$ . then  $U = (A + Bi)E$  and  $U^\dagger = (A - Bi)E$ , showing how the Fermion operators are expressed in terms of the simpler Clifford algebra of Majorana operators ( $A$  and  $B$  generating the split quaternions).

### 1.1. Writing in the Full Dirac Algebra

We have written the Dirac equation in one dimension of space and one dimension of time. We now boost the formalism directly to three dimensions of space. We take an independent Clifford algebra generated by  $\sigma_1, \sigma_2, \sigma_3$  with  $\sigma_i^2 = 1$  for  $i = 1, 2, 3$  and  $\sigma_i\sigma_j = -\sigma_j\sigma_i$  for  $i \neq j$ . Now assume that  $\alpha$  and  $\beta$  as we have used them above generate an independent Clifford algebra that commutes with the algebra of the  $\sigma_i$ . Replace the scalar momentum  $p$  by a 3-vector momentum  $p = (p_1, p_2, p_3)$  and let  $p \bullet \sigma = p_1\sigma_1 + p_2\sigma_2 + p_3\sigma_3$ . We replace  $\partial/\partial x$  with  $\nabla = (\partial/\partial x_1, \partial/\partial x_2, \partial/\partial x_3)$  and  $\partial p/\partial x$  with  $\nabla \bullet p$ .

We then have the following form of the Dirac equation.

$$i\partial\psi/\partial t = -i\alpha\nabla \bullet \sigma\psi + \beta m\psi.$$

Let  $\mathcal{O} = i\partial/\partial t + i\alpha\nabla \bullet \sigma - \beta m$  so that the Dirac equation takes the form  $\mathcal{O}\psi(x, t) = 0$ .

In analogy to our previous discussion we let  $\psi(x, t) = e^{i(p \bullet x - Et)}$  and construct solutions by first applying the Dirac operator to this  $\psi$ . The two Clifford algebras interact to generalize directly the nilpotent solutions and Fermion algebra, that we have detailed for one spatial dimension, to this three dimensional case. To this purpose the modified Dirac operator is

$$\mathcal{D} = i\beta\alpha\partial/\partial t + \beta\nabla \bullet \sigma - \alpha m.$$

And we have that  $\mathcal{D}\psi = U\psi$  where  $U = \beta\alpha E + \beta p \bullet \sigma - \alpha m$ . We have that  $U^2 = 0$  and  $U\psi$  is a solution to the modified Dirac Equation, just as before. And just as before, we can articulate the structure of the Fermion operators and locate the corresponding Majorana Fermion operators.

### 1.2. Majorana Fermions

There is more to do. We now discuss making Dirac algebra distinct from the one generated by  $\alpha, \beta, \sigma_1, \sigma_2, \sigma_3$  to obtain an equation that can have real solutions. This was the strategy that Majorana [3] followed to construct his Majorana Fermions. A real equation can have solutions that are invariant under complex conjugation and so can correspond to particles that are their own anti-particles. We will describe this Majorana algebra in terms of the split quaternions  $\epsilon$  and  $\eta$ . For convenience we use the matrix representation given below.

$$\epsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \eta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Let  $\hat{\epsilon}$  and  $\hat{\eta}$  generate another, independent algebra of split quaternions, commuting with the first algebra generated by  $\epsilon$  and  $\eta$ . Then a totally real Majorana Dirac equation can be written as follows:

$$(\partial/\partial t + \hat{\eta}\eta\partial/\partial x + \epsilon\partial/\partial y + \hat{\epsilon}\eta\partial/\partial z - \hat{\epsilon}\hat{\eta}\eta m)\psi = 0.$$

To see that this is a correct Dirac equation, note that

$$\hat{E} = \alpha_x \hat{p}_x + \alpha_y \hat{p}_y + \alpha_z \hat{p}_z + \beta m$$

(Here the ‘‘hats’’ denote the quantum differential operators corresponding to the energy and momentum.) will satisfy

$$\hat{E}^2 = \hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2 + m^2$$

if the algebra generated by  $\alpha_x, \alpha_y, \alpha_z, \beta$  has each generator of square one and each distinct pair of generators anti-commuting. From there we obtain the general Dirac equation by replacing  $\hat{E}$  by  $i\partial/\partial t$ , and  $\hat{p}_x$  with  $-i\partial/\partial x$  (and same for  $y, z$ ).

$$(i\partial/\partial t + i\alpha_x\partial/\partial x + i\alpha_y\partial/\partial y + i\alpha_z\partial/\partial z - \beta m)\psi = 0.$$

This is equivalent to

$$(\partial/\partial t + \alpha_x\partial/\partial x + \alpha_y\partial/\partial y + \alpha_z\partial/\partial z + i\beta m)\psi = 0.$$

Thus, here we take

$$\alpha_x = \hat{\eta}\eta, \alpha_y = \epsilon, \alpha_z = \hat{\epsilon}\eta, \beta = i\hat{\epsilon}\hat{\eta}\eta,$$

and observe that these elements satisfy the requirements for the Dirac algebra. Note how we have a significant interaction between the commuting square root of minus one ( $i$ ) and the element  $\hat{\epsilon}\hat{\eta}$  of square minus one in the split quaternions. This brings us back to considerations about the source of the square root of minus one. Both viewpoints combine in the element  $\beta = i\hat{\epsilon}\hat{\eta}\eta$  that makes this Majorana algebra work. Since the algebra appearing in the Majorana Dirac operator is constructed entirely from two commuting copies of the split quaternions, there is no appearance of the complex numbers, and when written out in  $2 \times 2$  matrices we obtain coupled real differential equations to be solved. This is a beginning of a new study of Majorana Fermions. For more information about this viewpoint, see [9]. In the next section we rewrite the Majorana Dirac operator, guided by nilpotents, obtaining solutions that directly use the Majorana Fermion operators.

## 2. Nilpotents, Majorana Fermions and the Majorana-Dirac Equation

Let  $\mathcal{D} = (\partial/\partial t + \hat{\eta}\eta\partial/\partial x + \epsilon\partial/\partial y + \hat{\epsilon}\eta\partial/\partial z - \hat{\epsilon}\hat{\eta}\eta m)$ . In the last section we have shown how  $\mathcal{D}$  can be taken as the Majorana operator through which we can look for real solutions to the Dirac equation. Letting  $\psi(x, t) = e^{i(p \bullet r - Et)}$ , we have

$$\mathcal{D}\psi = (-iE + i(\hat{\eta}\eta p_x + \epsilon p_y + \hat{\epsilon}\eta p_z) - \hat{\epsilon}\hat{\eta}\eta m)\psi.$$

Let

$$\Gamma = (-iE + i(\hat{\eta}\eta p_x + \epsilon p_y + \hat{\epsilon}\eta p_z) - \hat{\epsilon}\hat{\eta}\eta m)$$

and

$$U = \epsilon\eta\Gamma = (i(-\eta\epsilon E - \hat{\eta}\epsilon p_x + \eta p_y - \epsilon\hat{\epsilon}p_z) + \epsilon\hat{\epsilon}\hat{\eta}m).$$

The element  $U$  is nilpotent,  $U^2 = 0$ , and we have that  $U = A + iB$ ,  $AB + BA = 0$ ,  $A = \epsilon\hat{\epsilon}\hat{\eta}m$ ,  $B = -\eta\epsilon E - \hat{\eta}\epsilon p_x + \eta p_y - \epsilon\hat{\epsilon}p_z$ ,  $A^2 = -m^2$ , and  $B^2 = -E^2 + p_x^2 + p_y^2 + p_z^2 = -m^2$ .

Letting  $\nabla = \epsilon\eta\mathcal{D}$ , we have a new Majorana Dirac operator with  $\nabla\psi = U\psi$  so that  $\nabla(U\psi) = U^2\psi = 0$ . Letting  $\theta = (p \bullet r - Et)$ , we have  $U\psi = (A + Bi)e^{i\theta} = (A + Bi)(\text{Cos}(\theta) + i\text{Sin}(\theta)) = (A\text{Cos}(\theta) -$

$$BSin(\theta) + i(BCos(\theta) + ASin(\theta)).$$

Thus we have found two real solutions to the Majorana Dirac Equation:

$$\Phi = ACos(\theta) - BSin(\theta),$$

$$\Psi = BCos(\theta) + ASin(\theta)$$

with  $\theta = (p \bullet r - Et)$  and  $A$  and  $B$  the Majorana operators

$$A = \epsilon \hat{\epsilon} \hat{\eta} m,$$

$$B = -\eta \epsilon E - \hat{\eta} \epsilon p_x + \eta p_y - \epsilon \hat{\epsilon} p_z.$$

Note how the Majorana Fermion algebra generated by  $A$  and  $B$  comes into play in the construction of these solutions. This answers a natural question about the Majorana Fermion operators. Should one take the Majorana operators themselves seriously as representing physical states? Our calculation suggests that one should take them seriously.

In other work [4–7] we review the main features of recent applications of the Majorana algebra and its relationships with representations of the braid group and with topological quantum computing. The present analysis of the Majorana Dirac equation first appears in our paper [9].

### 3. Spacetime in 1+1 Dimensions

We begin this section by discussing an algebra that is directly related to Clifford algebra. As we shall see, this algebra is also inherent in the Dirac equation when we use light cone coordinates.

#### 3.1. Clifford algebra and Fermion algebra.

In two by two matrix algebra, we can take

$$\epsilon = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \eta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = a + b.$$

Here

$$a = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, b = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Thus

$$ab = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, ba = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

so that

$$a^2 = b^2 = 0,$$

$$a + b = \eta,$$

$$a - b = \epsilon \eta,$$

$$ab + ba = 1,$$

$$ab - ba = \epsilon.$$

More abstractly, suppose that we have a Clifford algebra generated by elements  $\epsilon$  and  $\eta$  with  $\epsilon^2 = \eta^2 = 1$  and  $\epsilon\eta + \eta\epsilon = 0$ . Then we can define new elements  $a$  and  $b$  by the equations

$$\begin{aligned}\eta &= a + b, \\ \epsilon\eta &= a - b.\end{aligned}$$

This means that

$$\begin{aligned}a &= \frac{1}{2}(1 + \epsilon)\eta, \\ b &= \frac{1}{2}(1 - \epsilon)\eta,\end{aligned}$$

from which it follows that

$$a^2 = b^2 = 0, ab + ba = 1.$$

Note that we are given that the starting Clifford algebra is associative and so further identities such as

$$aba = a, bab = b, abab = ab, baba = ba$$

follow easily from the given identities. We call an associative algebra generated by  $a, b$  with

$$a^2 = b^2 = 0, ab + ba = 1$$

a *Fermion algebra* since the annihilation, creation algebra for Fermions in quantum theory satisfies these identities. We see here that Clifford algebras (with an even number of generators) and Fermion algebras are interchangeable via the above transformations. This fact has been used by writers on Clifford algebras, [11] since it is useful to have projector properties such as  $(ab)(ab) = ab$ .

**Remark.** The above construction of Fermion algebra from Clifford algebra occurs without invoking an extra commuting square root of negative unity. It is common in physical applications to use a parallel construction involving  $i$  where  $i^2 = -1$  and  $i$  commutes with all elements of the algebra. One can then define  $\psi = \frac{1}{2}(\eta + i\epsilon)$  and  $\psi^\dagger = \frac{1}{2}(\eta - i\epsilon)$ . It follows that  $\psi^2 = (\psi^\dagger)^2 = 0$  and  $\psi\psi^\dagger + \psi^\dagger\psi = 1$ , and one has a Fermion algebra with complex conjugation constructed in relation to a Clifford algebra. Another relation with a commuting  $i$  occurs if we take

$$\begin{aligned}a &= (i/2)(\alpha\beta + \beta) \\ b &= (i/2)(\alpha\beta - \beta)\end{aligned}$$

where  $\alpha$  and  $\beta$  form a Clifford algebra with  $\alpha^2 = \beta^2 = 1$  and  $\alpha\beta + \beta\alpha = 0$ . Then  $a$  and  $b$  satisfy the Fermion relations and

$$\begin{aligned}ab + ba &= 1, \\ ab - ba &= \alpha,\end{aligned}$$

but

$$\begin{aligned}a + b &= i\alpha\beta, \\ a - b &= i\beta.\end{aligned}$$

Notice that  $(i\alpha\beta)^2 = +1$  while  $(i\beta)^2 = -1$ . Thus we can regard this as a re-writing of the previous pattern with

$$i\alpha\beta = \eta$$

and

$$i\beta = \epsilon\eta$$

so that

$$\alpha = \beta\beta\alpha = -\beta\alpha\beta = i\beta[\alpha\beta = \epsilon\eta\eta = \epsilon.$$

This means that this Fermion algebra can occur with or without the explicit commuting square root of negative unity,  $i$ .

### 3.2. The Dirac Equation in Light Cone Coordinates

Recall the translation of operators to light cone coordinate operators.

$$\hat{E} = i\partial/\partial t = (i/2)(\partial/\partial r + \partial/\partial l)$$

$$\hat{p} = (1/i)\partial/\partial x = (1/2i)(\partial/\partial r - \partial/\partial l)$$

Here is the nilpotent version of the Dirac operator as we have formulated it.

$$\mathcal{D} = \alpha\beta\hat{E} + \beta\hat{p} - \alpha\hat{m}$$

We translate this operator into light cone coordinates.

$$\mathcal{D} = \alpha\beta((i/2)(\partial/\partial r + \partial/\partial l)) + \beta((1/2i)(\partial/\partial r - \partial/\partial l)) - \alpha m$$

$$\mathcal{D} = i[(\alpha\beta + \beta)/2]\partial/\partial l + i[(\alpha\beta - \beta)/2]\partial/\partial r - \alpha m$$

Thus

$$\mathcal{D} = A\partial/\partial l + B\partial/\partial r - \alpha m$$

$$A = (i/2)(\alpha\beta + \beta)$$

$$B = (i/2)(\alpha\beta - \beta)$$

As the reader can see, we arrive at algebraic coefficients that we have described above as the Fermion algebra associated with the Clifford algebra generated by  $\alpha$  and  $\beta$ .

$$A + B = i\alpha\beta$$

$$A - B = i\beta$$

Further relations take the form:

$$AB + BA = 1, AB - BA = \alpha, A^2 = B^2 = 0, \alpha^2 = 1$$

$$A\alpha = -A, \alpha A = A, B\alpha = B, \alpha B = -B$$

Thus

$$A\alpha + \alpha A = 0, B\alpha + \alpha B = 0$$

$$A\beta + \beta A = i, B\beta + \beta B = -i$$

Let

$$\psi = e^{i(rX - lY)}$$

where

$$X = p - E$$

and

$$Y = p + E.$$

Thus

$$XY = -m^2.$$

Then

$$\mathcal{D}\psi = U\psi$$

where

$$U = -iAX + iBY - \alpha m.$$

Thus

$$U^2 = ABXY + BAXY + m^2 = XY + m^2 = p^2 - E^2 + m^2 = 0.$$

Note that with

$$U^\dagger = -iAY + iBX - \alpha m$$

we have

$$(U^\dagger)^2 = 0$$

and

$$UU^\dagger + U^\dagger U = 4E^2$$

### 3.3. Solving the 1+1 Dirac Equation

The (real valued) Majorana version of the Dirac operator

$$\mathcal{D} = A\partial/\partial l + B\partial/\partial r - \alpha m$$

that we have discussed above can be taken via the representation

$$A = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \alpha = \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Then

$$AB = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, BA = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, AB - BA = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = \alpha.$$

Letting

$$\Theta = rX - lY,$$

and

$$S = \text{Sin}(\Theta), C = \text{Cos}(\Theta),$$

we have

$$U\psi = U(C + iS) = (AXS - BYS - \alpha mC) + i(-AXC + BYC - \alpha mS).$$

In the matrix representation we find

$$AXS - BYS - \alpha mC = \begin{pmatrix} mC & -YS \\ XS & -mC \end{pmatrix}.$$

And from this, letting

$$\psi_1 = mC, \psi_2 = XS$$

we have

$$\partial\psi_1/\partial r = -mXS = -m\psi_2$$

and

$$\partial\psi_2/\partial l = -XYC = m^2C = m\psi_1$$

Thus

$$\partial\psi_1/\partial r = -m\psi_2$$

$$\partial\psi_2/\partial l = m\psi_1$$

Note that these equations are satisfied by

$$\psi_1 = -m\text{Sin}(-(E-p)r - (E+p)l),$$

$$\psi_2 = (E+p)\text{Cos}(-(E-p)r - (E+p)l)$$



exactly when  $E^2 = p^2 + m^2$  as we have assumed. It is quite interesting to see these direct solutions to the Dirac equation emerge in this 1 + 1 case. The solutions are fundamental and they are distinct from the usual solutions that emerge from the Feynman checkerboard Model [1,2]. It is the above equations that form the basis for the Feynman checkerboard model that is obtained by examining paths in a discrete Minkowski plane generating a path integral for the Dirac equation.

**Remark.** Note that a simplest instance of the above form of solution is obtained by writing

$$e^{i(r+l)} = \cos(r+l) + i\sin(r+l) = \sum_{n=0}^{\infty} (\sqrt{-1})^n \sum_{i+j=n} \frac{r^i l^j}{i! j!}.$$

Then with  $\psi_2 = \cos(r+l)$  and  $\psi_1 = \sin(r+l)$  we have  $\partial\psi_1/\partial l = \psi_2$ ,  $\partial\psi_2/\partial r = -\psi_1$ , solving the Dirac equation in the case where  $m = 1$ .

**Remark.** Let  $\psi_R = \sum_{k=0}^{\infty} (-1)^k \frac{r^{k+1} l^k}{(k+1)! k!}$ ,  $\psi_L = \sum_{k=0}^{\infty} (-1)^k \frac{r^k l^{k+1}}{k! (k+1)!}$ ,  $\psi_0 = \sum_{k=0}^{\infty} (-1)^k \frac{r^k l^k}{k! k!}$ . Then  $\psi_1 = \psi_0 + \psi_L$  and  $\psi_2 = \psi_0 - \psi_R$  give a solution to the Dirac equation in light cone coordinates as we have written it above with  $m = 1$ :  $\partial\psi_1/\partial l = \psi_2$ ,  $\partial\psi_2/\partial r = -\psi_1$ . These series are shown in [2] to be a natural limit of evaluations of sums of discrete paths on the Feynman checkerboard. The key to our earlier approach is that if one defines

$$C[\Delta]_k^x = \frac{(x)(x-\Delta)(x-2\Delta)\cdots(x-(k-1)\Delta)}{k!},$$

Then  $C[\Delta]_k^x$  takes the role of  $\frac{x^k}{k!}$  for discrete different derivatives with step length  $\Delta$  and it can be interpreted as a choice coefficient. A Feynman path on a rectangle in Minkowski space can be interpreted as two choice of  $k$  or  $k+1$  points along the  $r$  and  $l$  edges of the rectangle. Thus the products in the limit expressions of the form  $\frac{r^k l^{k+1}}{k! (k+1)!}$  or  $\frac{r^{k+1} l^k}{(k+1)! k!}$  correspond to paths on the checkerboard with  $k$  corners in a limit where there are infinitely many such paths. The details are in our paper [2]. The solutions we have given above, motivated by the Majorana algebra, are related in form to these path sum solutions. Our solutions contain more information, related to the factorization  $(E-p)(E+p) = E^2 - p^2 = m^2$ . In the usual checkerboard solution the propagators only know about the mass and not its factorization relative to energy and momentum. More work needs to be done to fully understand the relationship of solutions to the Dirac equation and path summations.

## Список литературы/References

1. R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, 1965, McGraw Hill Companies, Inc, New York.
2. L. H. Kauffman and H. P. Noyes [1996], Discrete Physics and the Dirac Equation, *Physics Letters A*, 218 ,pp. 139-146.
3. E. Majorana [1937], A symmetric theory of electrons and positrons, *I Nuovo Cimento*, **14** (1937), pp. 171-184.
4. L. H. Kauffman [1991,1994,2001,2012], Knots and Physics, *World Scientific Pub.*, Singapore.
5. L. H. Kauffman [2016], Knot logic and topological quantum computing with Majorana Fermions. In "Logic and algebraic structures in quantum computing and information Lecture Notes in Logic, J. Chubb, J. Chubb, Ali Eskandarian, and V. Harizanov, editors, 124 pages Cambridge University Press (2016).
6. L. H. Kauffman [2017], Iterants, *Entropy* (2017), 19, 347; doi:10.3390/e19070347.
7. L. H. Kauffman [2018], Majorana Fermions and representations of the braid group. *Internat. J. Modern Phys. A*, 33 (2018), no. 23, 1830023, 28 pp.
8. L. H. Kauffman and S. J. Lomonaco Jr. [2007],  $q$ -deformed spin networks, knot polynomials and anyonic topological quantum computation. *J. Knot Theory Ramifications*, 16 (2007), no. 3, 267-332.

9. L. H. Kauffman and P. Rowlands [2021], The Dirac equation and the Majorana Dirac equation, *Aporia - 40-th Anniversary ANPA Proceedings - 1979-2019*, Edited by John Ceres Amson, ANPA Pub. February 2021.
10. P. Rowlands [2007], Zero to Infinity - The Foundations of Physics, *Series on Knots and Everything*, vol. 41, World Scientific Publishing Co., 2007.
11. G. Sobczyk, *Matrix Gateway to Geometric Algebra, Spacetime and Spinors*, Independent Publisher, Amazon-KDP (2019).

### Авторы

**Кауффман Луис Хирш**, профессор, Факультет математики, статистики и вычислительной техники, Иллинойский университет в Чикаго, 851 Саут Морган Стрит, Чикаго, Иллинойс, 60607-7045, США.

E-mail: kauffman@uic.edu

### Просьба ссылаться на эту статью следующим образом:

Кауффман Л. Х. Примечания к уравнению Дирака. *Пространство, время и фундаментальные взаимодействия*. 2022. № 40. С. 64–73.

### Authors

**Kauffman Louis Hirsch**, professor, Department of Mathematics, Statistics and Computer Science, University of Illinois at Chicago, 851 South Morgan Street, Chicago, Illinois, 60607-7045, USA.

E-mail: kauffman@uic.edu

### Please cite this article in English as:

Kauffman L. H. Remarks on the Dirac Equation. *Space, Time and Fundamental Interactions*, 2022, no. 40, pp. 64–73.